



Lot Specific Information

Lot Number:	Lot Specific*
Volume:	Lot Specific*
Concentration:	Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein:	Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	BE0063
Clone:	DTA-1
Isotype:	Rat IgG2b, λ
Recommended Isotype Control(s):	InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	Mouse CD25+ CD4+ T cells
Reported Applications:	<i>in vivo</i> GITR stimulation
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	<2EU/mg (<0.002EU/ μ g) Determined by LAL gel clotting assay
Purity:	>95% Determined by SDS-PAGE
Sterility:	0.2 μ m filtered
Production:	Purified from tissue culture supernatant in an animal free facility
Purification:	Protein G
RRID:	AB_1107688
Molecular Weight:	150 kDa

Description

The DTA-1 monoclonal antibody reacts with mouse GITR (glucocorticoid-induced TNFR-related gene), a 66-70 kDa co-stimulatory immune checkpoint molecule belonging to the Tumor Necrosis Factor superfamily (TNFRSF18). GITR is expressed at low levels on resting T lymphocytes and at high levels on regulatory T cells. GITR is upregulated on activated T cells where it provides co-stimulation. GITR ligand (GITRL) is found on B cells, macrophages, dendritic and endothelial cells, and is implicated in regulating both innate and adaptive immune responses. GITR is also thought to play a key role in dominant immunological self-tolerance maintained by regulatory T cells. Knockout studies in mice also suggest the role of this receptor is in the regulation of CD3-driven T cell activation and programmed cell death. The DTA-1 antibody is an agonistic antibody that is commonly used to induce GITR signaling *in vivo*.

Shelf-life and Storage

Store at the stock concentration at 4°C. **Do not freeze.**
All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at bxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit <https://bxcell.com/product/m-gitr/#references> or scan the QR code below.

Bio X Cell, Inc.

bxcell.com
1.866.787.3444
customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell



Bio X Cell, Inc.

bxcell.com
1.866.787.3444

customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell