



**Lot Specific Information**

<b>Lot Number:</b>	Lot Specific*
<b>Volume:</b>	Lot Specific*
<b>Concentration:</b>	Lot Specific* (generally 4 to 11 mg/ml) *
<b>Total Protein:</b>	Lot Specific*

\*This information will be noted on the certificate of analysis that ships with this product.

**Product Information**

<b>Catalog Number:</b>	<b>BE0075-1</b>
<b>Clone:</b>	<b>1A8</b>
<b>Isotype:</b>	Rat IgG2a, κ
<b>Recommended Isotype Control(s):</b>	InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
<b>Recommended Dilution Buffer:</b>	InVivoPure pH 7.0 Dilution Buffer
<b>Immunogen:</b>	EL4J cells transfected with Ly6G <i>in vivo</i> neutrophil depletion <i>in vivo</i> MDSC depletion Immunofluorescence Immunohistochemistry (paraffin) Immunohistochemistry (frozen) Flow cytometry
<b>Reported Applications:</b>	
<b>Formulation:</b>	PBS, pH 7.0 Contains no stabilizers or preservatives
<b>Endotoxin:</b>	<2EU/mg (<0.002EU/μg) Determined by LAL gel clotting assay
<b>Purity:</b>	>95% Determined by SDS-PAGE
<b>Sterility:</b>	0.2 μM filtered
<b>Production:</b>	Purified from tissue culture supernatant in an animal free facility
<b>Purification:</b>	Protein G
<b>RRID:</b>	AB_1107721
<b>Molecular Weight:</b>	150 kDa

**Description**

The 1A8 monoclonal antibody reacts with mouse Ly6G. Ly6G is a 21-25 kDa member of the Ly-6 superfamily of GPI-anchored cell surface proteins with roles in cell signaling and cell adhesion. Ly6G is expressed differentially during development by cells in the myeloid lineage including monocytes, macrophages, granulocytes, and neutrophils. Monocytes typically express Ly6G transiently during development while mature granulocytes and peripheral neutrophils retain expression making Ly6G a good cell surface marker for these populations. Unlike the RB6-8C5 antibody, the 1A8 antibody reacts specifically with mouse Ly6G with no reported cross reactivity with Ly6C.

**Shelf-life and Storage**

Store at the stock concentration at 4°C. **Do not freeze.**  
All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at [bxcell.com/faqs](https://bxcell.com/faqs).

**Protocol Information**

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

**Application References**

For a complete list of references, visit <https://bxcell.com/product/invivomab-anti-m-ly-6g/#references> or scan the QR code below.

**Bio X Cell, Inc.**

bxcell.com  
1.866.787.3444  
[customerservice@bxcell.com](mailto:customerservice@bxcell.com)

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.  
Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell



---

**Bio X Cell, Inc.**

bxcell.com  
1.866.787.3444

**customerservice@bxcell.com**

*Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.*

**Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell**