

## InVivoMAb anti-human/rat/fish AChR

### Lot Specific Information

<b>Lot Number:</b>	Lot Specific*
<b>Volume:</b>	Lot Specific*
<b>Concentration:</b>	Lot Specific* (generally 4 to 11 mg/ml) *
<b>Total Protein:</b>	Lot Specific*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

<b>Catalog Number:</b>	<b>BE0123</b>
<b>Clone:</b>	<b>Mab35 (TIB-175)</b>
<b>Isotype:</b>	Rat IgG1
<b>Recommended Isotype Control(s):</b>	InVivoMAb rat IgG1 isotype control, anti-horseradish peroxidase
<b>Recommended Dilution Buffer:</b>	InVivoPure pH 7.0 Dilution Buffer
<b>Immunogen:</b>	<i>Electrophorus electricus</i> acetylcholine receptor
<b>Reported Applications:</b>	EAMG induction in rats Immunofluorescence
<b>Formulation:</b>	PBS, pH 7.0 Contains no stabilizers or preservatives
<b>Endotoxin:</b>	<2EU/mg (<0.002EU/μg) Determined by LAL gel clotting assay
<b>Purity:</b>	>95% Determined by SDS-PAGE
<b>Sterility:</b>	0.2 μM filtered
<b>Production:</b>	Purified from tissue culture supernatant in an animal free facility
<b>Purification:</b>	Protein G
<b>RRID:</b>	AB_10950153
<b>Molecular Weight:</b>	150 kDa

### Description

The mAb35 (TIB-175) antibody reacts with human rat and fish acetylcholine receptor (AChR). The AChR is a heterodimeric receptor consisting of four subunits in a molar stoichiometry  $\alpha 2\beta\gamma\delta$  during the early embryonic stages or after denervation and  $\alpha 2\beta\epsilon\delta$  in the adult form. The mAb35 antibody binds to the major immunogenic region of the  $\alpha$  subunits. It has been shown to passively transfer experimental autoimmune myasthenia gravis.

### Shelf-life and Storage

Store at the stock concentration at 4°C. **Do not freeze.**

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at [bxcell.com/faqs](https://bxcell.com/faqs).

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

### Application References

For a complete list of references, visit <https://bxcell.com/product/anti-achr/#references> or scan the QR code below.



---

**Bio X Cell, Inc.**

bxcell.com  
1.866.787.3444

**customerservice@bxcell.com**

*Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.*

**Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell**