Technical Data Sheet

InVivoMAb anti-mouse MHC Class I (H-2Kd, H-2Dd)



Lot Specific Information

Lot Number:Lot Specific*Volume:Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:BE0180Clone:34-1-2SIsotype:Mouse IgG2a, K

Recommended Isotype Control(s): InVivoMAb mouse IgG2a isotype control, unknown specificity

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer

Immunogen: BDF mouse spleen cells

Reported Applications: in vivo activation of APCs

Formulation: PBS, pH 7.0

Contains no stabilizers or preservatives

Endotoxin: <2EU/mg (<0.002EU/µg)
Determined by LAL gel clotting assay

Purity: Determined by SDS-PAGE

Sterility: 0.2 µM filtered

Production: Purified from tissue culture supernatant in an animal free facility

Purification:Protein GRRID:AB_10950841Molecular Weight:150 kDa

Description

The 34-1-2S monoclonal antibody is reported to react with the mouse $H-2K^b$ and $H-2D^d$ MHC class I alloantigens. MHC class I antigens are heterodimers consisting of one alpha chain (44 kDa) associated with β 2 microglobulin (11.5 kDa). The antigen is expressed by all nucleated cells at varying levels. MHC Class I molecules present endogenously synthesized antigenic peptides to CD8 T cells.

Shelf-life and Storage

Store at the stock concentration at $4\,^{\circ}\text{C}.$ Do not freeze.

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at bxcell.com/fags.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bxcell.com/product/h-2-k-d-d-d/#references or scan the QR code below.

Bio X Cell, Inc.

bxcell.com

1.866.787.3444

customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.



Bio X Cell, Inc.

bxcell.com 1.866.787.3444

customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.