

InVivoMAb anti-mouse/human/rat CCL2 (MCP-1)

Lot Specific Information

Lot Number:	Lot Specific*
Volume:	Lot Specific*
Concentration:	Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein:	Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	BE0185
Clone:	2H5
Isotype:	Armenian Hamster IgG, κ
Recommended Isotype Control(s):	InVivoMAb polyclonal Armenian hamster IgG
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	CHO-expressed mouse MCP-1 <i>in vivo</i> CCL2 neutralization Immunohistochemistry (frozen)
Reported Applications:	PBS, pH 7.0 Contains no stabilizers or preservatives
Formulation:	<2EU/mg (<0.002EU/μg) Determined by LAL gel clotting assay
Endotoxin:	>95% Determined by SDS-PAGE
Purity:	0.2 μM filtered
Sterility:	Purified from tissue culture supernatant in an animal free facility
Production:	Protein G
Purification:	AB_10950302
RRID:	150 kDa
Molecular Weight:	

Description

The 2H5 antibody reacts with mouse, human, and rat CCL2 also known as monocyte chemoattractant protein 1 (MCP-1). CCL2 is a 13 kDa chemokine that is produced mainly by monocytes/macrophages but is also expressed by endothelial, fibroblasts, epithelial, smooth muscle, mesangial, astrocytic, and microglial cells. It can be expressed constitutively or after induction by oxidative stress, cytokines, or growth factors. CCL2 regulates the migration and infiltration of monocytes, memory T lymphocytes, and NK cells and is implicated in the pathogenesis of several diseases characterized by monocytic infiltrates, such as multiple sclerosis, psoriasis, rheumatoid arthritis, and atherosclerosis. The 2H5 antibody has been shown to neutralize the biological effects of CCL2 *in vivo*.

Shelf-life and Storage

Store at the stock concentration at 4°C. **Do not freeze.**

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at bxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit <https://bxcell.com/product/m-r-h-ccl2-mcp-1/#references> or scan the QR code below.

Bio X Cell, Inc.

bxcell.com
1.866.787.3444
customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell



Bio X Cell, Inc.

bxcell.com

1.866.787.3444

customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell