



**Lot Specific Information**

<b>Lot Number:</b>	Lot Specific*
<b>Volume:</b>	Lot Specific*
<b>Concentration:</b>	Lot Specific* (generally 4 to 11 mg/ml) *
<b>Total Protein:</b>	Lot Specific*

\*This information will be noted on the certificate of analysis that ships with this product.

**Product Information**

<b>Catalog Number:</b>	<b>BE0261</b>
<b>Clone:</b>	<b>KT3</b>
<b>Isotype:</b>	Rat IgG2a
<b>Recommended Isotype Control(s):</b>	InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
<b>Recommended Dilution Buffer:</b>	InVivoPure pH 7.0 Dilution Buffer
<b>Immunogen:</b>	CBAT6 mouse thymocytes <i>in vitro</i> T cell negative selection <i>in vitro</i> T cell stimulation/activation Immunofluorescence
<b>Reported Applications:</b>	
<b>Formulation:</b>	PBS, pH 7.0 Contains no stabilizers or preservatives
<b>Endotoxin:</b>	<2EU/mg (<0.002EU/μg) Determined by LAL gel clotting assay
<b>Purity:</b>	>95% Determined by SDS-PAGE
<b>Sterility:</b>	0.2 μM filtered
<b>Production:</b>	Purified from tissue culture supernatant in an animal free facility
<b>Purification:</b>	Protein G
<b>RRID:</b>	AB_2687740
<b>Molecular Weight:</b>	150 kDa

**Description**

The KT3 monoclonal antibody reacts with mouse CD3ε, a 20 kDa transmembrane cell-surface protein that belongs to the immunoglobulin superfamily. CD3ε is one of five polypeptide chains that combine to form the TCR complex. CD3ε is expressed on T lymphocytes, NK-T cells, and to varying degrees on developing thymocytes. CD3 plays roles in TCR signaling, T lymphocyte activation, and antigen recognition. The KT3 antibody has been shown to induce T lymphocyte activation via binding and stimulating the TCR.

**Shelf-life and Storage**

Store at the stock concentration at 4°C. **Do not freeze.**  
All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at [bxcell.com/faqs](https://bxcell.com/faqs).

**Protocol Information**

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

**Application References**

For a complete list of references, visit <https://bxcell.com/product/invivomab-cd3e/#references> or scan the QR code below.

**Bio X Cell, Inc.**

bxcell.com  
1.866.787.3444  
[customerservice@bxcell.com](mailto:customerservice@bxcell.com)

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.  
Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell



---

**Bio X Cell, Inc.**

bxcell.com  
1.866.787.3444

**customerservice@bxcell.com**

*Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.*

**Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell**