

## InVivoPlus anti-mouse TIM-3 (CD366)

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bxcell.com/terms-and-conditions/>.

### Lot Specific Information

<b>Lot Number:</b>	Lot Specific*
<b>Volume:</b>	Lot Specific*
<b>Concentration:</b>	Lot Specific* (generally 4 to 11 mg/ml) *
<b>Total Protein:</b>	Lot Specific*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

<b>Catalog Number:</b>	<b>BP0115</b>
<b>Clone:</b>	<b>RMT3-23</b>
<b>Isotype:</b>	Rat IgG2a, κ
<b>Recommended Isotype Control(s):</b>	InVivoPlus rat IgG2a isotype control, anti-trinitrophenol
<b>Recommended Dilution Buffer:</b>	InVivoPure pH 7.0 Dilution Buffer
<b>Immunogen:</b>	Recombinant mouse TIM-3 <i>in vivo</i> TIM-3 neutralization <i>in vitro</i> TIM-3 blocking Flow cytometry
<b>Reported Applications:</b>	PBS, pH 7.0 Contains no stabilizers or preservatives
<b>Formulation:</b>	<1EU/mg (<0.001EU/μg) Determined by LAL gel clotting assay
<b>Endotoxin:</b>	>95% Determined by SDS-PAGE
<b>Purity:</b>	0.2 μM filtered
<b>Sterility:</b>	Purified from tissue culture supernatant in an animal free facility
<b>Production:</b>	Protein G
<b>Purification:</b>	<5% Determined by DLS
<b>Aggregation:</b>	AB_10949464
<b>RRID:</b>	150 kDa
<b>Molecular Weight:</b>	

### Murine Pathogen Test Results

Mouse Norovirus: Negative, Mouse Parvovirus: Negative, Mouse Minute Virus: Negative, Mouse Hepatitis Virus: Negative, Reovirus Screen: Negative, Lymphocytic Choriomeningitis virus: Negative, Lactate Dehydrogenase-Elevating Virus: Negative, Mouse Rotavirus: Negative, Theiler's Murine Encephalomyelitis: Negative, Ectromelia/Mousepox Virus: Negative, Hantavirus: Negative, Polyoma Virus: Negative, Mouse Adenovirus: Negative, Sendai Virus: Negative, Mycoplasma Pulmonis: Negative, Pneumonia Virus of Mice: Negative, Mouse Cytomegalovirus: Negative, K Virus: Negative,

### Description

The RMT3-23 monoclonal antibody reacts with mouse TIM-3 (T cell immunoglobulin and mucin domain-3) also known as CD366. TIM-3 is a 60 kDa member of the TIM family of immune checkpoint receptors and exists as a type I transmembrane glycoprotein with a mucin-like domain in its extracellular portion and a tyrosine phosphorylation motif in its cytoplasmic portion. TIM-3 is specifically expressed at high levels on the surface of Th1 lymphocytes whereas Th2 lymphocytes express TIM-1 and TIM-2. TIM-3 activation occurs via binding to the cell-associated C-type lectin galectin-9. Upon binding TIM-3 induces apoptosis of Th1 cells. Inhibition of TIM-3 signaling in mice has been shown to exacerbate experimental autoimmune encephalomyelitis, promote IFNγ production and Th1 cell proliferation. Tim-3 has also been shown to be required for the induction of tolerance, as both TIM-3 knockout animals and mice treated with TIM-3-Ig fusion protein display defects in the induction of antigen-specific tolerance. Additionally, TIM-3 signaling is currently being explored as a cancer immunotherapy target as CD8 T cells which express both TIM-3 and PD-1 exhibit greater defects in both cell-cycle progression and effector cytokine production than cells that express PD-1 alone. The RMT3-23 antibody acts as a TIM-3 receptor antagonist and has been shown to have functional activity including suppressing tumor cell growth in a murine sarcoma model.

### Shelf-life and Storage

#### Bio X Cell, Inc.

bxcell.com  
1.866.787.3444  
customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell

Store at the stock concentration at 4°C. **Do not freeze.**

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at [bxc.com/faq](https://bxc.com/faq).

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

### Application References

For a complete list of references, visit <https://bxc.com/product/invivoplus-anti-m-tim-3/#references> or scan the QR code below.



---

### Bio X Cell, Inc.

[bxc.com](https://bxc.com)  
1.866.787.3444

[customerservice@bxc.com](mailto:customerservice@bxc.com)

*Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.*

**Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell**